



# Nematiron Plus

Fertilizer

## Radicular bioestimulant

### Composition:

Free aminoacid: 6,1% w/w  
Glycine: 2,1% w/w, Lysine: 2% w/w, Glutamic acid: 2% w/w  
Total nitrogen: 1,0 % w/w, Organic nitrogen: 1,0% w/w

### Characterystics:

– Aspect: Yellow liquid – pH: 2,8 – Specific Gravity (20°C): 1,12 gr/cc

Liquid organic compound ready soluble in water. It is based on polycarboxilic acids and aminoacids obtained by natural fermentation process. It works as a powerful biostimulant and revitalizing agent of plant metabolism in stressful and adverse conditions, as droughts, intense cold temperature and nematode attack. It revitalizes the root system protecting it against stressful conditions improving plant defenses.

### Crops:

It is recommended for all type of crops.

### Instructions:

- \* Indicated to be used by fertigation as a source of organic matter:
  - In case of plant transplantsand inn soils with a high number of nematodes.
  - In any case where a bio-stimulating effect on the plant is required.
- \* Preventive application to improve and strengthen plant defenses against the nematode attack and other stressful conditions. Apply the product 5 to 10 days before sowing or crop transplant at a dose of 100 lt. Ha, irrigate previously with plenty of water during 2 to 3 hours and apply the product at the end of this irrigation period.

### Dosage:

Applications	Doses and Applications
STRAWBERRIES	Before crop: 60 - 75 l/Ha 3 Applications of 20 - 25 l/Ha. Firs one 15 days after transplant, the rest with 15 days in between until the end of the culture.
VEGETABLES/ ORNAMENTAL	Before crop: 60 - 75 l/Ha 1 application of 20 – 25 l/Ha every 15 days after transplanting up to 3 applications
BANANA (MORE THAN 1 YEAR OLD)	Before crop: 90 l/Ha After the first year: 3 applications of 30 l/Ha split within the year.
POTATOES AND SUGAR BEET	Before crop: 60 - 75 l/Ha 1 application of 25 l/Ha every 20 days beginning 15 days after bourgeoning, up to 3 applications.

### Packing:

Presentation in plastic bottles / jerrycans of 1,5,20 liter and IBC containers of 1000 liters.

It is recommended not to exceed the recommended doses

Note: The information contained in this sheet has been prepared according to our current knowledge and should only be understood as a guide. This company is solely responsible for the composition, formulation and content of its products and, in no case for the form of use made by the customer. The results and effectiveness of our products on a surface not exceeding 100m2. This company will not be, in any case responsible for the damages that may be caused in case of non-compliance with these instructions.